

A TORRES CONDITION FOR TWISTED ALEXANDER POLYNOMIALS

TAKAYUKI MORIFUJI

ABSTRACT. As a generalization of a fundamental result about the Alexander polynomial of links, we give a description of a Torres condition for the twisted Alexander polynomial of links associated to a unimodular representation.

1. INTRODUCTION

The theory of twisted Alexander polynomial was introduced by Lin [11] and Wada [15] independently. Lin defined it for knots in the 3-sphere using regular Seifert surfaces. On the other hand, Wada defined the twisted Alexander polynomial for finitely presentable groups, which include the link groups. In particular, as an application, Wada told the Kinoshita-Terasaka knot from the Conway knot by means of his invariant. Shortly afterward, several significant results on the original Alexander polynomial were generalized to the twisted case. For example, equivalence of the twisted Alexander polynomial and the Reidemeister torsion, and its symmetry [8], [6], sliceness obstruction for knots and a relation to the Casson-Gordon invariant [6], [7], monicness of the twisted Alexander polynomial for fibered knots [1], [2] and so on. Recently the twisted Alexander polynomials are extensively investigated. See for instance [3], [4], [5], [9], [10], [12] and [13].

However, almost all results mentioned above are basically about knots in the 3-sphere and it seems that there are few generalized results on links. The purpose of the present paper is to give a generalization of the following well-known formula for the Alexander polynomial of links.

Theorem 1.1 (Torres [14]). *For the Alexander polynomial $\Delta_L(t_1, \dots, t_\mu)$ of a μ -component link $L = L_1 \cup \dots \cup L_\mu$, it holds that*

$$\Delta_L(t_1, \dots, t_{\mu-1}, 1) = \begin{cases} \frac{t_1^{l_1} - 1}{t_1 - 1} \Delta_{L'}(t_1) & \text{if } \mu = 2 \\ (t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}} - 1) \Delta_{L'}(t_1, \dots, t_{\mu-1}) & \text{if } \mu > 2, \end{cases}$$

where $L' = L_1 \cup \dots \cup L_{\mu-1}$ is the link obtained from L by removing L_μ and l_i denotes the linking number of the components L_i and L_μ .

More precisely, we give a description of a Torres condition for the twisted Alexander polynomial of links associated to a unimodular representation. In the next section, we briefly recall the definition of the twisted Alexander polynomial for the link group. The precise statement and the proof of the main theorem of this paper are given in Section 3.

2000 *Mathematics Subject Classification.* Primary 57M25, Secondary 57M05, 57M27.
Key words and phrases. Torres condition, link group, twisted Alexander polynomial.

2. TWISTED ALEXANDER POLYNOMIAL FOR LINKS

Let $L = L_1 \cup \cdots \cup L_\mu$ be a μ -component link in the 3-sphere. We denote the fundamental group of its exterior E by $G(L)$. Namely, we put $G(L) = \pi_1(E)$ and call it the link group. We choose and fix a Wirtinger presentation of $G(L)$:

$$G(L) = \langle x_1, \dots, x_u \mid r_1, \dots, r_{u-1} \rangle.$$

Then the abelianization homomorphism

$$\alpha : G(L) \rightarrow H_1(E; \mathbb{Z}) \cong \mathbb{Z}^{\oplus \mu} = \langle t_1 \rangle \oplus \cdots \oplus \langle t_\mu \rangle$$

is given by assigning to each generator x_i the meridian element $t_k \in H_1(E; \mathbb{Z})$ of the corresponding component L_k of L . In this paper, we consider a linear representation $\rho : G(L) \rightarrow SL(n; F)$, where F denotes a field.

These maps naturally induce two ring homomorphisms $\tilde{\rho} : \mathbb{Z}[G(L)] \rightarrow M(n; F)$ and $\tilde{\alpha} : \mathbb{Z}[G(L)] \rightarrow \mathbb{Z}[t_1^{\pm 1}, \dots, t_\mu^{\pm 1}]$, where $\mathbb{Z}[G(L)]$ is the group ring of $G(L)$ over $\mathbb{Z}$ and $M(n; F)$ is the matrix algebra of degree n over F . Taking the tensor of $\tilde{\rho}$ and $\tilde{\alpha}$, we obtain a ring homomorphism

$$\tilde{\rho} \otimes \tilde{\alpha} : \mathbb{Z}[G(L)] \rightarrow M(n; F[t_1^{\pm 1}, \dots, t_\mu^{\pm 1}]).$$

Let F_u denote the free group on generators $x_1, \dots, x_u$ and

$$\Phi : \mathbb{Z}[F_u] \rightarrow M(n; F[t_1^{\pm 1}, \dots, t_\mu^{\pm 1}])$$

the composite of the surjection $\mathbb{Z}[F_u] \rightarrow \mathbb{Z}[G(L)]$ induced by the presentation and the map $\tilde{\rho} \otimes \tilde{\alpha}$.

Let us consider the $(u-1) \times u$ matrix $M = M(t_1, \dots, t_\mu)$ whose (i, j) th component is the $n \times n$ matrix

$$\Phi \left(\frac{\partial r_i}{\partial x_j} \right) \in M(n; F[t_1^{\pm 1}, \dots, t_\mu^{\pm 1}]),$$

where $\partial/\partial x$ denotes the free differential calculus. This matrix M is called the Alexander matrix of $G(L)$ associated to the representation ρ .

For $1 \leq j \leq u$, let us denote by $M_j = M_j(t_1, \dots, t_\mu)$ the $(u-1) \times (u-1)$ matrix obtained from M by removing the column corresponding to a generator x_j . We also regard M_j as an $n(u-1) \times n(u-1)$ matrix with coefficients in $F[t_1^{\pm 1}, \dots, t_\mu^{\pm 1}]$.

Then Wada's twisted Alexander polynomial of a link L for a representation $\rho : G(L) \rightarrow SL(n; F)$ is defined to be a rational function

$$\Delta_{L, \rho}(t_1, \dots, t_\mu) = \frac{|M_j|}{|\Phi(x_j - 1)|},$$

where $|M_j|$ denotes the determinant of the matrix M_j , and it is well-defined up to a factor $\pm t_1^{nk_1} \cdots t_\mu^{nk_\mu}$ ($k_i \in \mathbb{Z}$) if n is odd and up to only $t_1^{nk_1} \cdots t_\mu^{nk_\mu}$ if n is even (see [15] Section 5 for details).

Remark 2.1. In general, the twisted Alexander polynomial for finitely presentable groups is a rational function, but it is actually a polynomial for the link groups (see [15] Proposition 9 and [9] Theorem 3.1).

3. A TORRES CONDITION

In this section, we state and prove a generalized Torres condition for the twisted Alexander polynomial of links. An advantage of our description here is that we need not separate the case for $\mu = 2$ from $\mu > 2$. We first prove the theorem in the case of an $SL(2; F)$ -representation. After reading the proof for it, one can easily show the similar result for general cases.

Theorem 3.1. *Let $L = L_1 \cup \dots \cup L_\mu$ be a μ -component link and $L' = L_1 \cup \dots \cup L_{\mu-1}$. For a given representation $\rho' : G(L') \rightarrow SL(2; F)$, it holds that*

$$\Delta_{L, \rho}(t_1, \dots, t_{\mu-1}, 1) = \{(t_1^{l_1} \dots t_{\mu-1}^{l_{\mu-1}})^2 - \varepsilon_{\rho'} t_1^{l_1} \dots t_{\mu-1}^{l_{\mu-1}} + 1\} \Delta_{L', \rho'}(t_1, \dots, t_{\mu-1}),$$

where $\rho : G(L) \rightarrow SL(2; F)$ is the composite of the natural surjection $G(L) \rightarrow G(L')$ and ρ' , l_i denotes the linking number of L_i and L_μ , and $\varepsilon_{\rho'}$ is an element of F .

Proof. For the link group $G(L)$, we choose a Wirtinger presentation:

$$G(L) = \langle x_{ij} \mid r_{kl} \rangle,$$

where $x_{i1}, x_{i2}, \dots, x_{ij_i}$ ($1 \leq i \leq \mu$) are generators corresponding to the component L_i and the relation

$$r_{kl} = x_{k'l'}^{\pm 1} x_{kl} x_{k'l'}^{\mp 1} x_{k,l+1}^{-1}$$

corresponds to a crossing of $L_{k'}$ over L_k . We should note that the link group $G(L)$ has the deficiency one.

Let us consider the Alexander matrix of $G(L)$ associated to the representation $\rho : G(L) \rightarrow SL(2; F)$:

$$M(t_1, \dots, t_\mu) = \left(\Phi \left(\frac{\partial r_{kl}}{\partial x_{ij}} \right) \right).$$

Then we know that if we remove the column corresponding to a generator x_{ij} ,

$$|M_{ij}(t_1, \dots, t_\mu)| = |\Phi(x_{ij} - 1)| \Delta_{L, \rho}(t_1, \dots, t_\mu)$$

holds. Thus if we set $t_\mu = 1$ in $M(t_1, \dots, t_\mu)$, it follows that

$$|M_{ij}(t_1, \dots, t_{\mu-1}, 1)| = |\Phi(x_{ij} - 1)| \Delta_{L, \rho}(t_1, \dots, t_{\mu-1}, 1)$$

if $i \neq \mu$.

Now the generators $\{x_{\mu j}\}$ appear in the following two kinds of relations:

$$(i) \ r_{\mu j} = x_{rs}^{\pm 1} x_{\mu j} x_{rs}^{\mp 1} x_{\mu, j+1}^{-1} \quad \text{and} \quad (ii) \ r_{pq} = x_{\mu l}^{\pm 1} x_{pq} x_{\mu l}^{\mp 1} x_{p, q+1}^{-1},$$

where the relation (i) corresponds to crossings of L_r over L_μ and (ii) corresponds to that of L_μ over L_p . Let us see which are the contributions of these relations to the matrix $M(t_1, \dots, t_{\mu-1}, 1)$. First the contributions of $r_{\mu j}$ are as follows:

$$\begin{aligned} \Phi \left(\frac{\partial r_{\mu j}}{\partial x_{rs}} \right)_{t_\mu=1} &= O, \\ \Phi \left(\frac{\partial r_{\mu j}}{\partial x_{\mu j}} \right)_{t_\mu=1} &= \begin{cases} t_r^{\pm 1} \rho(x_{rs})^{\pm 1} & \text{if } \mu \neq r \\ I & \text{if } \mu = r, \end{cases} \\ \Phi \left(\frac{\partial r_{\mu j}}{\partial x_{\mu, j+1}} \right)_{t_\mu=1} &= -I, \end{aligned}$$

where O and I denote the zero and the identity matrix respectively. We have used here the fact that $\rho(x_{\mu j}) = I$ for $1 \leq j \leq j_\mu$ (because the generators $\{x_{\mu j}\}$ are in

the kernel of the natural surjective homomorphism from $G(L)$ to $G(L')$). Next the contributions of r_{pq} are as follows:

$$\begin{aligned}\Phi\left(\frac{\partial r_{pq}}{\partial x_{\mu l}}\right)_{t_{\mu}=1} &= \pm(I - t_p \rho(x_{pq})), \\ \Phi\left(\frac{\partial r_{pq}}{\partial x_{pq}}\right)_{t_{\mu}=1} &= I, \\ \Phi\left(\frac{\partial r_{pq}}{\partial x_{p,q+1}}\right)_{t_{\mu}=1} &= -\rho(x_{pq})\rho(x_{p,q+1})^{-1} \text{ if } p \neq \mu,\end{aligned}$$

and the case $p = \mu$ has already been considered. Therefore we see that the matrix $M(t_1, \dots, t_{\mu-1}, 1)$ has the following form:

$$M(t_1, \dots, t_{\mu-1}, 1) = \begin{pmatrix} A & B \\ O & C \end{pmatrix},$$

where

$$A = \left(\Phi\left(\frac{\partial r_{kl}}{\partial x_{ij}}\right)_{t_{\mu}=1} \right) (k, i \neq \mu), \quad B = \left(\Phi\left(\frac{\partial r_{kl}}{\partial x_{\mu j}}\right)_{t_{\mu}=1} \right) (k \neq \mu, 1 \leq j \leq j_{\mu}),$$

and

$$\begin{aligned}C &= \left(\Phi\left(\frac{\partial r_{\mu l}}{\partial x_{\mu j}}\right)_{t_{\mu}=1} \right) (1 \leq j, l \leq j_{\mu}) \\ &= \begin{pmatrix} t_{r_1}^{\delta_{r_1}} \rho(x_{r_1 s_1})^{\delta_{r_1}} & & & -I & & \\ & t_{r_2}^{\delta_{r_2}} \rho(x_{r_2 s_2})^{\delta_{r_2}} & & & -I & \\ & & t_{r_3}^{\delta_{r_3}} \rho(x_{r_3 s_3})^{\delta_{r_3}} & & & \\ & & & \ddots & & \\ & & & & -I & \\ -I & & & & & t_{r_{j_{\mu}}}^{\delta_{r_{j_{\mu}}}} \rho(x_{r_{j_{\mu}} s_{j_{\mu}}})^{\delta_{r_{j_{\mu}}}} \end{pmatrix}.\end{aligned}$$

In the submatrix C , there is an appearance of $t_i^{\delta_i}$ for each crossing of L_i over L_{μ} ($1 \leq i \leq \mu$) and $\delta_i = 1$ or -1 according as L_i crosses over L_{μ} from left to right or from right to left. Thereby, we obtain

$$\begin{aligned}|C| &= \prod_{i=1}^{j_{\mu}} |\rho(x_{r_i s_i})^{\delta_{r_i}}| t_{r_i}^{2\delta_{r_i}} - \sum_{\sigma \in S} c_1^{\sigma} \cdots c_{j_{\mu}}^{\sigma} t_{r_1}^{\delta_{r_1}} \cdots t_{r_{j_{\mu}}}^{\delta_{r_{j_{\mu}}}} + 1 \\ &= (t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}})^2 - \varepsilon_{\rho'} t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}} + 1,\end{aligned}$$

where $c_i^{\sigma} \in F$ denotes a component of the matrix $\rho(x_{r_i s_i})^{\delta_{r_i}}$ determined by a permutation σ and S is a subset of the symmetric group $\mathfrak{S}_{2j_{\mu}}$ consisting of permutations which choose just one component from each submatrix $\rho(x_{r_i s_i})^{\delta_{r_i}}$. Furthermore the submatrix A is equivalent to the Alexander matrix $M'(t_1, \dots, t_{\mu-1})$ of $G(L')$ associated to the representation $\rho' : G(L') \rightarrow SL(2; F)$. Hence if we remove a column corresponding to a generator x_{ij} ($i \neq \mu$), then we have

$$\begin{aligned}|M_{ij}(t_1, \dots, t_{\mu-1}, 1)| &= |A_{ij}| |C| \\ &= \{(t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}})^2 - \varepsilon_{\rho'} t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}} + 1\} |M'_{ij}(t_1, \dots, t_{\mu-1})|,\end{aligned}$$

where A_{ij} is the matrix obtained from A by removing the column corresponding to x_{ij} . Therefore, by definition of the twisted Alexander polynomial, we see that

$$\Delta_{L,\rho}(t_1, \dots, t_{\mu-1}, 1) = \{(t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}})^2 - \varepsilon_{\rho'} t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}} + 1\} \Delta_{L',\rho'}(t_1, \dots, t_{\mu-1}).$$

This completes the proof of Theorem 3.1. $\square$

Remark 3.2. The fact that $\Delta_{L,\rho}(t_1, \dots, t_{\mu-1}, 1)$ is divisible by $\Delta_{L',\rho'}(t_1, \dots, t_{\mu-1})$ also follows from a recent result of Kitano, Suzuki and Wada in [10]. However, we can have no detailed information on the quotient from their result.

A linear representation $\rho : G(L) \rightarrow GL(n; F)$ is called reducible if it has a non-trivial invariant subspace in F^n . In this case, we can obtain a piece of information about the coefficient $\varepsilon_{\rho'}$.

Corollary 3.3. *Under the setting as in Theorem 3.1, if $\rho' : G(L') \rightarrow SL(2; F)$ is a reducible representation, then we have*

$$\varepsilon_{\rho'} = \lambda^l + \lambda^{-l} \quad (l = l_1 + \cdots + l_{\mu-1}),$$

where λ is an eigenvalue of the image of a generator of $G(L')$.

Proof. First we can assume that the images of generators in a Wirtinger presentation of $G(L)$ have the following forms:

$$\rho(x_{ij}) = \begin{pmatrix} a_{ij} & b_{ij} \\ 0 & a_{ij}^{-1} \end{pmatrix} \quad (i \neq \mu) \quad \text{and} \quad \rho(x_{\mu j}) = I,$$

where $a_{ij} \in F^\times$ and $b_{ij} \in F$. Because the representation ρ' has a 1-dimensional invariant subspace in F^2 .

Since $x_{ij}x_{kl}^{-1}$ ($i, k \neq \mu$) is an element of the commutator subgroup $[G(L), G(L)]$, we see that $a_{ij} = a_{kl}$ holds for these generators. We then put $\lambda = a_{ij}$ for simplicity. Each lower left component of $\rho(x_{ij})$ is zero, so that the nontrivial terms appeared in the coefficient of $t_{r_1}^{\delta_{r_1}} \cdots t_{r_{j\mu}}^{\delta_{r_{j\mu}}}$ are just

$$-\lambda^{\delta_{r_1} + \cdots + \delta_{r_{j\mu}}} - \lambda^{-(\delta_{r_1} + \cdots + \delta_{r_{j\mu}})} = -(\lambda^l + \lambda^{-l}),$$

where $l = \sum l_i$. This completes the proof. $\square$

Example 3.4. Let $\rho' : G(L') \rightarrow SL(2; F)$ be a reducible representation of the knot $L' = L_1$. Then the twisted Alexander polynomial of L' associated to ρ' is given by

$$\Delta_{L',\rho'}(t_1) = \frac{\Delta_{L'}(\lambda t_1) \Delta_{L'}(\lambda^{-1} t_1)}{(t_1 - \lambda)(t_1 - \lambda^{-1})},$$

where $\Delta_{L'}(t_1)$ is the original Alexander polynomial of L' (see the proof of [9] Theorem 3.1 for instance). Hence we have

$$\begin{aligned} \Delta_{L,\rho}(1, 1) &= \frac{(1 - \lambda^{l_1})(1 - \lambda^{-l_1})}{(1 - \lambda)(1 - \lambda^{-1})} \Delta_{L'}(\lambda) \Delta_{L'}(\lambda^{-1}) \\ &= (1 + \lambda + \cdots + \lambda^{l_1-1})(1 + \lambda^{-1} + \cdots + \lambda^{-(l_1-1)}) \Delta_{L'}(\lambda) \Delta_{L'}(\lambda^{-1}). \end{aligned}$$

In particular, if ρ' is trivial (namely, $\lambda = 1$), then we obtain $\Delta_{L,\rho}(1, 1) = l_1^2$ (because $\Delta_{L'}(1) = \pm 1$).

Example 3.5. Let $\rho' : G(L') \rightarrow SL(2; F)$ be the trivial representation. In this case $\varepsilon_{\rho'} = 2$ holds, so that we have

$$\Delta_{L,\rho}(t_1, \dots, t_{\mu-1}, 1) = (t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}} - 1)^2 \Delta_{L',\rho'}(t_1, \dots, t_{\mu-1}).$$

This formula corresponds to the square of Torres' original formula in Theorem 1.1. In particular, $\Delta_{L,\rho}(1, \dots, 1) = 0$ holds for $\mu > 2$.

If we slightly modify the proof of Theorem 3.1, we obtain the following general formula for a unimodular representation $\rho' : G(L') \rightarrow SL(n; F)$. We omit here the repetitious proof.

Theorem 3.6. Let $L = L_1 \cup \cdots \cup L_\mu$ be a μ -component link and $L' = L_1 \cup \cdots \cup L_{\mu-1}$. For a given representation $\rho' : G(L') \rightarrow SL(n; F)$, it holds that

$$\begin{aligned} \Delta_{L,\rho}(t_1, \dots, t_{\mu-1}, 1) = & \{(t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}})^n + \sum_{k=1}^{n-1} \varepsilon_{k,\rho'} (t_1^{l_1} \cdots t_{\mu-1}^{l_{\mu-1}})^{n-k} + (-1)^n\} \\ & \times \Delta_{L',\rho'}(t_1, \dots, t_{\mu-1}), \end{aligned}$$

where $\rho : G(L) \rightarrow SL(n; F)$ is the composite of the natural surjection $G(L) \rightarrow G(L')$ and ρ' , l_i denotes the linking number of L_i and L_μ , and $\varepsilon_{k,\rho'}$ ($1 \leq k \leq n-1$) are elements of F .

Finally, we extend Corollary 3.3 when all the images of the representation $\rho' : G(L') \rightarrow SL(n; F)$ are upper triangle matrices.

Corollary 3.7. Under the setting as in Theorem 3.6, if $\text{Im}(\rho')$ are upper triangle matrices, then the coefficient $\varepsilon_{k,\rho'}$ is given by

$$\varepsilon_{k,\rho'} = (-1)^k \sum_{1 \leq i_1 < \cdots < i_k \leq n} (\lambda_1 \cdots \hat{\lambda}_{i_1} \cdots \hat{\lambda}_{i_k} \cdots \lambda_n)^l,$$

where λ_k ($1 \leq k \leq n$) are the eigenvalues of the image of a generator of $G(L')$ and $\hat{\lambda}_k$ implies that λ_k is removed from the product.

Acknowledgements. This paper was written while the author was visiting the Ludwig-Maximilians-Universität in München. He would like to express his sincere thanks for their hospitality.

REFERENCES

1. J. C. Cha, *Fibred knots and twisted Alexander invariants*, Trans. Amer. Math. Soc. **355** (2003), 4187–4200.
2. H. Goda, T. Kitano and T. Morifuji, *Reidemeister torsion, twisted Alexander polynomial and fibered knots*, Comment. Math. Helv. **80** (2005), 51–61.
3. H. Goda and T. Morifuji, *Twisted Alexander polynomial for $SL(2, \mathbb{C})$ -representations and fibered knots*, C. R. Math. Acad. Sci. Soc. R. Can. **25** (2003), 97–101.
4. M. Heusener and J. Porti, *Deformations of reducible representations of 3-manifold groups into $PSL(2, \mathbb{C})$* , math.GT/0411365.
5. B. Jiang and S. Wang, *Twisted topological invariants associated with representations*, in Topics in Knot Theory (1993), 211–227.
6. P. Kirk and C. Livingston, *Twisted Alexander invariants, Reidemeister torsion, and Casson-Gordon invariants*, Topology **38** (1999), 635–661.
7. P. Kirk and C. Livingston, *Twisted knot polynomials: inversion, mutation and concordance*, Topology **38** (1999), 663–671.
8. T. Kitano, *Twisted Alexander polynomial and Reidemeister torsion*, Pacific J. Math. **174** (1996), 431–442.

9. T. Kitano and T. Morifuji, *Divisibility of twisted Alexander polynomial and fibered knots*, preprint.
10. T. Kitano, M. Suzuki and M. Wada, *Twisted Alexander polynomial and surjectivity of a group homomorphism*, preprint.
11. X. S. Lin, *Representations of knot groups and twisted Alexander polynomials*, Acta Math. Sin. (Engl. Ser.) **17** (2001), 361–380.
12. T. Morifuji, *Twisted Alexander polynomial for the braid group*, Bull. Austral. Math. Soc. **64** (2001), 1–13.
13. M. Suzuki, *Twisted Alexander polynomial for the Lawrence-Krammer representation*, Bull. Austral. Math. Soc. **70** (2004), 67–71.
14. G. Torres, *On the Alexander polynomial*, Ann. of Math. **57** (1953), 57–89.
15. M. Wada, *Twisted Alexander polynomial for finitely presentable groups*, Topology **33** (1994), 241–256.

DEPARTMENT OF MATHEMATICS, TOKYO UNIVERSITY OF AGRICULTURE AND TECHNOLOGY, KOGANEI, TOKYO 184-8588, JAPAN

E-mail address: morifuji@cc.tuat.ac.jp